

Abnormal Measurements	Consensus 2012	Supporting References	Notes
Nuchal Fold 15- 18wk	≥5mm	Gray 1994	
Nuchal Fold 18-22 wks	≥6mm	Gray 1994	
Cisterna Magna	10mm	Smith 1986 & Twickler 2002	
Lateral Ventricles	10mm	Daddario 2012 & Farrell 1994	
Choroid separation	≥4mm	Hertzberg 1994 & Bronsteen 2006	
Cavum Septum Pellucidum	Chart	Serhatlioglu 2003 & Falco 2000	
Cerebellum	GA mm	Smith 1986	
Pericardial Effusion	>3mm	Brown 1998	
Bladder (Early 1st trimester)	7mm	Liao 2003	
Bladder (2nd trimester)	Chart	Maizels 2004	
Kidneys- Renal Pelviectasis			If Caliectasis present may use lower cutoff value for follow up
Early 2nd trimester (13-16 wks)	4mm	Nguyen 2010	
2nd trimester (17-32 wks)	4mm	Nguyen 2010	
3rd trimester (32-40 wks)	7mm	Nguyen 2010	
F/u for pyelectasis	32		
Low Lying Placenta	≥2.5cm	Creasy & Resnik	≥2.5cm no need for FU or CS in patient without bleeding.
Marginal Previa	At os	Creasy & Resnik	
Marginal Placental Cord Insertion	2cm	Liu 2002	

Cervix	<3cm	SOGC guideline 2011	<3cm at 24 weeks on TAUS should trigger TVUS
Uterine Artery Dopplers P.I.	1.6		
EFW % age that needs dopplers	10%	SMFM guideline 2012	
Oligohydramnios			
Single vertical pocket	2cm	Chauhan 2004	
AFI	5cm	Chauhan 2004	
Polyhydramnios			
Single vertical pocket	>8cm		
AFI	>25cm		
F/u for velamentous PCI	28wk	Sepulveda 2003	
F/u for placenta previa	28wk	SOGC guideline 2007	
F/u for vasa previa	28wk	SOGC guideline 2009	
Multiples			
Oligo - single pocket	<2cm		
Poly - single pocket	>8cm		